

Appendix to “Inconsistency of Inaccessibility”

Alexander Kiselev

aakiselev@yahoo.com

This paper is the concise addition to the foregoing work “Inconsistency of Inaccessibility”, containing the presentation of main theorem proof (in ZF) about inaccessible cardinals nonexistence. Here some refinement of this presentation is set forth. Much attention is devoted to the explicit and substantial development and cultivation of basic ideas, serving as grounds for all main constructions and reasonings.

The proof of main theorem in its brief form is exposed in specified work [1], one can also see it at arXiv site [1]. The present paper is its short complement using previous notions and designations from [1]. The proof of main theorem has been published in 1997, its systematic exposition has been published in 2000, and the most complete and detailed form the proof of inaccessible cardinals nonexistence have received in works “Inaccessibility and Subinaccessibility”, Part I [2] and Part II [3] in 2008, 2010; these two works one can see also at arXiv sites [2], [3].

Remind, that further matrics and matrix functions reduced to χ^* are considered, the lower index χ^* will be omitted in notations, for instance $\alpha_{\chi^*}^\downarrow$, $\alpha_{\chi^*}^\downarrow$, S_{χ^*f} , $S_{\chi^*\tau}$ will be denoted through $\alpha^\downarrow$, $\alpha^\downarrow$, S_f , S_τ and so on.

Here it remains only to eliminate the case when $\delta = \chi^{*+}$, that is when the function $S_f^{<\alpha^0}$ is $\leq$ -nondecreasing up to χ^{*+} . It means that the unrelativized function S_f , being $\leq$ -bounded by the fixed ordinal $Od(S_{\tau^*}) < \chi^{*+}$, loses this boundedness after its relativization to α^0 , that is to its transformation into $S_f^{<\alpha^0}$. One can see that it happens because of deformation of the universe after its relativization to α^0 : because of losing properties of subinaccessibility of all levels $\geq n$ by all cardinals $\leq \gamma_{\tau_0}$ after their relativization to α^0 (Kiselev [4]).

So, many important properties of lower levels of the universe do not extend up to relativizing cardinals, namely to prejump cardinals of reduced matrices that are values of matrix functions. And all the rest part of this work consists in correction of this lack. In order to prevent this phenomenon we introduce special cardinals named disseminators that are extending such properties without distortion:

Definition 1 Let $0 < \alpha < \alpha_1 \leq k$, $X \subseteq \alpha$, $X \neq \emptyset$.

The ordinal α is named disseminator of level m with data base X below α_1 iff for every $\mathbf{n} \in \omega_0$ and train $\vec{a}$ of ordinals $\in X$ there holds $U_m^{\Sigma < \alpha_1}(\mathbf{n}, \vec{a}) \longrightarrow U_m^{\Sigma < \alpha}(\mathbf{n}, \vec{a})$. The formula defining in L_k the set of all such disseminators is denoted by $SIN_m^{<\alpha_1}[X](\alpha)$; this set itself is also denoted by $SIN_m^{<\alpha_1}[X]$, while its disseminators – by the common symbol δ^X or, briefly, by δ , pointing to α_1 , X in the context. Everywhere further $m=n+1$ will be considered. ⊥

Remind, the symbol U_{n+1}^Σ here denotes the Σ_{n+1} -formula which is universal for the class of Σ_{n+1} -formulas but without any occurrences of the constant $\underline{l}$. The term “disseminator”

is justified by the fact that such an ordinal really extends Π_{n+1} -properties (containing constants from its base) from lower levels of the universe up to relativizing cardinal α_1 .

The discussion of disseminators is more clear when data base X is some ordinal ρ (that is the set of all smaller ordinals). In this case we shall write $SIN_{n+1}^{<\alpha_1} [<\rho]$ instead of $SIN_{n+1}^{<\alpha_1} [X]$; further ρ is closed under the pair function. Now one should supply singular matrixes with disseminators:

Definition 2 Let $\chi^* < \alpha < \alpha_1 \leq k$ and S be a matrix reduced to cardinal χ^* and singular on a carrier α .

1) We name as disseminator for S on α (or as disseminator for this carrier) of the level $n+1$ with data base ρ every disseminator $\delta \in SIN_{n+1}^{<\alpha^\downarrow} [<\rho] \cap SIN_n^{<\alpha^\downarrow}$ below the prejump cardinal $\alpha^\downarrow$ after χ^* .

2) The matrix S is called the disseminator matrix or, more briefly, the δ -matrix of the level $n+1$ admissible on the carrier α for $\gamma = \gamma_\tau^{<\alpha_1}$ below α_1 iff it possesses some disseminator $\delta < \gamma$ of the level $n+1$ with some base $\rho \leq \chi^{*+}$ on this carrier such that $S \triangleleft \rho$ (also below α_1). $\dashv$

Now we are going to undertake the second approach to the main theorem proof idea. Remind that the simplest matrix function S_f possess the property of monotonicity, but the direct proof of the required contradiction – the proof of its nonmonotonicity – is hampered by the occasion: some essential properties of lower levels of universe do not extend up to prejump cardinals of matrices on their carriers that are values of matrix functions.

In order to destroy this obstacle we equipped such matrices with corresponding disseminators and as the result the simplest matrix functions are transformed to their more complicated forms, that is to δ -functions:

Definition 3 Let $\gamma < \alpha < \alpha_1 \leq k$. 1) We denote through $\mathbf{K}_n^{\forall < \alpha_1}(\gamma, \alpha)$ the formula:

$$SIN_{n-1}^{<\alpha_1}(\gamma) \wedge \forall \gamma' \leq \gamma (SIN_n^{<\alpha_1}(\gamma') \longrightarrow SIN_n^{<\alpha}(\gamma')) .$$

If this formula is fulfilled by constants γ, α, α_1 , then we say that α conserves $SIN_n^{<\alpha_1}$ -cardinals $\leq \gamma$ below α_1 . If S is a matrix on a carrier α and its prejump cardinal $\alpha^\downarrow$ after χ^* conserves these cardinals, then we also say that S on α conserves these cardinals below α_1 .

2) We denote through $\mathbf{K}_{n+1}^\exists(\chi^*, \delta, \gamma, \alpha, \rho, S)$ the Π_{n-2} -formula:

$$\sigma(\chi^*, \alpha, S) \wedge \chi^* < \delta < \gamma < \alpha \wedge S \triangleleft \rho \leq \chi^{*+} \wedge SIN_{n+1}^{<\alpha^\downarrow} [<\rho](\delta) \wedge SIN_n^{<\alpha^\downarrow}(\delta).$$

Here, remind, the formula $\sigma(\chi^*, \alpha, S)$ means that S is singular matrix on its carrier α reduced to the cardinal χ^* ; δ is disseminator for S on α with the base ρ of the level $n+1$. We denote through $\mathbf{K}^{<\alpha_1}(\chi^*, \delta, \gamma, \alpha, \rho, S)$ the formula:

$$\mathbf{K}_n^{\forall < \alpha_1}(\gamma, \alpha^\downarrow) \wedge \mathbf{K}_{n+1}^{\exists \triangleleft \alpha_1}(\chi^*, \delta, \gamma, \alpha, \rho, S) \wedge \alpha < \alpha_1 .$$

3) If this formula is fulfilled by constants $\chi^*, \delta, \gamma, \alpha, \rho, S, \alpha_1$, then we say that $\chi^*, \delta, \alpha, \rho, S$ are strongly admissible for γ below α_1 . If some of them are fixed or meant by the context, then we say that others are strongly admissible for them (and for γ) below α_1 .

4) The matrix S is called strongly disseminator matrix or, briefly, δ -matrix strongly admissible on the carrier α for $\gamma = \gamma_\tau^{<\alpha_1}$ below α_1 , iff it possesses some disseminator $\delta < \gamma$ with base ρ strongly admissible for them (also below α_1). In every case of this kind δ -matrix is denoted by the common symbol δS or S . $\dashv$

Definition 4 Let $\chi^* < \alpha_1$. We call as the matrix δ -function of the level n below α_1 reduced to χ^* the function $\delta S_f^{<\alpha_1} = (\delta S_\tau^{<\alpha_1})_\tau$ taking values for $\tau < k$:

$$\delta S_{\tau}^{<\alpha_1} = \min_{\leq} \{ S \triangleleft \chi^{*+} : \exists \delta, \alpha, \rho < \gamma_{\tau+1}^{<\alpha_1} \mathbf{K}^{<\alpha_1}(\chi^*, \delta, \gamma_{\tau}^{<\alpha_1}, \alpha, \rho, S) \}. \quad \dashv$$

Therefore such function has values that are singular strongly admissible on their carriers δ -matrices.

The unrelativized function δS_f really does exist on the final subinterval of the inaccessible cardinal k as it shows

Lemma 5 (About δ -function definiteness) *There exists an ordinal $\delta < k$ such that δS_f is defined on the set $T = \{\tau : \delta < \gamma_{\tau} < k\}$.*

The minimal of such ordinals δ is denoted by δ^* and its index by τ_1^* , so that $\delta^* = \gamma_{\tau_1^*}$ $\dashv$

The monotonicity of this function is treated in the previous way: the function $\delta S_{\tau}^{<\alpha_1}$ is called (totally) monotone iff $\forall \tau', \tau'' (\tau_1^* < \tau' < \tau'' < k \longrightarrow \delta S_{\tau'} \leq \delta S_{\tau''})$.

Let's discuss the situation which arises. For some convenience the suitable notation of ordinal intervals will be used for $\alpha_1 < \alpha_2$: $[\alpha_1, \alpha_2[= \alpha_2 - \alpha_1$; $] \alpha_1, \alpha_2[= \alpha_2 - (\alpha_1 + 1)$ and so on (here ordinals α_1, α_2 are sets of smaller ordinals).

It was noted above, that the simplest matrix function S_f is $\leq$ -monotone, but for every $\tau > \tau^*$ the prejump cardinal $\alpha^{\downarrow}$ of S_{τ} on its corresponding carrier $\alpha \in]\gamma_{\tau}, k[$ do not conserve the subinaccessibility of levels $\geq n$ of cardinals $\leq \gamma_{\tau}$, and some other important properties of the lower levels of the universe are destroyed when relativizing to $\alpha^{\downarrow}$.

In order to overcome these obstruction we have supplied the values of this function, matrices δS_{τ} , by disseminators of the level $n+1$ and required the conservation of the subinaccessibility of the level n for cardinals $\leq \gamma_{\tau}$, that is we passed to the δ -function δS_f . And these disseminators by definitions 2, 3 really do overcome these obstructions and now basic Π_{n+1} -properties and all Π_n -properties of the lower levels of the universe are extending up to prejump cardinals of matrixes δS_{τ} carriers after χ^* .

But now it involves the new complication: now one can see, that after this modification such δ -function loses its property of (total) monotonicity on $[\tau_1^*, k[$, and just due to the fact that prejump cardinals $\alpha^{\downarrow}$ of δ -matrices carriers α , vice versa, generate the subinaccessibility of the level n of some cardinals $\leq \gamma_{\tau}$ that become subinaccessible (relatively to $\alpha^{\downarrow}$), not being those in the universe (Kiselev [5]).

So, here we come to the third and final approach to the main theorem proof idea:

The following requirements should be imposed on δ -matrices δS_{τ} successively in the course of matrix function defining:

- 1) *they must possess the property of “autoexorcizivity”, that is of self-exclusion in situations when $S = \delta S_{\tau}$ is the first value of δ -function monotonicity violation in its preceding part; the matrices with this property (of “unit characteristic”) will have the priority over other matrices (of “zero characteristic” respectively) in the course of defining the matrix function;*
- 2) *more requirements should be imposed on matrices of zero characteristic, preventing their forming: their disseminator data bases must increase substantially, and must be even disabled (“suppressed”), when the preceding part of matrix function, which has already been defined, contains monotonicity violation, in order to correct this fault – the using of matrices of zero characteristic;*

on this grounds δ -matrix functions should receive inconsistent properties of monotonicity and nonmonotonicity simultaneously.

Obviously, all these considerations require the recursive definition of matrix functions, setting there values depending on specified properties of their preceding values.

This resulting recursive definition follows definition 4, but in recursive form and the formula $\mathbf{K}^{<\alpha_1}$ transforms in its corresponding version which we denote through $\alpha\mathbf{K}^{* < \alpha_1}$. As the result there arises matrix functions which we denote through $\alpha S_f^{<\alpha_1}$ and name α -functions below α_1 .

One can see that these functions possess many basic properties of simplest matrix functions and δ -functions, for instance, analogous to their definiteness properties for unrelativized α -function αS_f : there exists the cardinal $\delta < k$ such that $\{\tau : \delta < \gamma_\tau < k\} \subseteq \text{dom}(\alpha S_f)$; more precisely:

Lemma 6 (About α -function definitness) *There exist cardinals $\delta < \gamma < k$ such that for every SIN_n -cardinal $\alpha_1 > \gamma$, $\alpha_1 < k$ limit for $SIN_n \cap \alpha_1$ the function $\alpha S_f^{<\alpha_1}$ is defined on the nonempty set $T^{\alpha_1} = \{\tau : \delta < \gamma_\tau^{<\alpha_1} < \alpha_1\}$.*

The minimal of such cardinals δ is denoted by $\alpha\delta^*$ and its index by $\alpha\tau_1^* = \tau(\alpha\delta^*)$, so that $\alpha\delta^* = \gamma_{\alpha\tau_1^*}$. $\dashv$

Now the monotonicity of α -function is considered; here this notion is used again in the previous sense: the function $\alpha S_f^{<\alpha_1}$ is called monotone on interval $[\tau_1, \tau_2[$ below α_1 iff $]\tau_1, \tau_2[\subseteq \text{dom}(\alpha S_f^{<\alpha_1})$ and $\forall \tau', \tau'' (\tau_1 < \tau' < \tau'' < \tau_2 \longrightarrow \alpha S_{\tau'}^{<\alpha_1} \leq \alpha S_{\tau''}^{<\alpha_1})$. As we shall see, this property is rather strong; in particular, any interval $[\tau_1, \tau_2[$ of its monotonicity can not be “too long” – that is the corresponding interval $]\gamma_{\tau_1}, \gamma_{\tau_2}[$ can not contain any SIN_n -cardinals; therefore $\alpha S_f^{<\alpha_1}$ doesn't possess this property in total sense:

Theorem 1. *Let $\alpha S_f^{<\alpha_1}$ be monotone on $[\tau_1, \tau_2[$ below α_1 and $\tau_1 = \min\{\tau :]\tau, \tau_2[\subseteq \text{dom}(\alpha S_f^{<\alpha_1})\}$, then $]\gamma_{\tau_1}^{<\alpha_1}, \gamma_{\tau_2}^{<\alpha_1}[\cap SIN_n^{<\alpha_1} = \emptyset$.* $\dashv$

The contradiction, which proves the main theorem, is the following:

On one hand, by lemma 6 function $\alpha S_f^{<\alpha_1}$ is defined on nonempty set $T^{\alpha_1} = \{\tau : \alpha\delta^* < \gamma_\tau < \alpha_1\}$ for every sufficiently great cardinal $\alpha_1 \in SIN_n$. Its monotonicity on this set is excluded by theorem 1.

But on the other hand, this monotonicity is ensured by the following theorem for every SIN_n -cardinal $\alpha_1 > \alpha\delta^*$ of sufficiently great cofinality:

Theorem 2. *Let function $\alpha S_f^{<\alpha_1}$ be defined on nonempty set $T^{\alpha_1} = \{\tau : \gamma_{\tau_1}^{<\alpha_1} < \gamma_\tau^{<\alpha_1} < \alpha_1\}$ for $\alpha_1 < k$ such that (i) $\tau_1 = \min\{\tau : \forall \tau' (\gamma_\tau^{<\alpha_1} < \gamma_{\tau'}^{<\alpha_1} \longrightarrow \tau' \in \text{dom}(\alpha S_f^{<\alpha_1}))\}$, (ii) $\sup SIN_n^{<\alpha_1} = \alpha_1$; (iii) $cf(\alpha_1) \geq \chi^{*+}$.*

Then $\alpha S_f^{<\alpha_1}$ is monotone on this set below α_1 : $\forall \tau_1, \tau_2 \in T^{\alpha_1} (\tau_1 < \tau_2 \rightarrow \alpha S_{\tau_1}^{<\alpha_1} \leq \alpha S_{\tau_2}^{<\alpha_1})$.

Proof is the following in general. The reasoning will be carried out by the induction on the cardinal α_1 . Suppose, that this theorem fails and the cardinal α_1^* is *minimal* breaking this theorem, that is the function $\alpha S_f^{<\alpha_1^*}$ is nonmonotone on the nonempty set $T^{\alpha_1^*} = \{\tau : \gamma_{\tau_1^*}^{<\alpha_1^*} < \gamma_\tau^{<\alpha_1^*} < \alpha_1^*\}$ with specified properties (i)–(iii) for some τ_1^* , so that the *inductive hypothesis* is accepted:

for every $\alpha_1 < \alpha_1^*$ the function $\alpha S_f^{<\alpha_1}$ is monotone on the nonempty set T^{α_1} with properties (i)–(iii).

It follows straight from theorem 1, that this α_1^* is simply the *minimal* cardinal α_1 , providing the existence of such nonempty set T^{α_1} because for every such $\alpha_1 < \alpha_1^*$ the function $\alpha S_f^{<\alpha_1}$ on T^{α_1} is *nonmonotone* by theorem 1 and at the same time is *monotone* by the minimality of α_1^* .

But in this situation all possible violations of function $\alpha S_f^{<\alpha_1^*}$ monotonicity on specified set $T^{\alpha_1^*}$ draws the existence of some nonempty set $T^{\alpha_1} = \{\tau : \gamma_{\tau_1}^{<\alpha_1} < \gamma_\tau^{<\alpha_1} < \alpha_1\}$ with

properties (i)–(iii) for lesser $\alpha_1 < \alpha_1^*$. So, the function $\alpha S_f^{<\alpha_1^*}$, nevertheless, is monotone on the set $T^{\alpha_1^*}$. $\dashv$

Now let us sum up. All the reasoning was conducted in the system

$$ZF + \exists k \text{ (} k \text{ is weakly inaccessible cardinal),}$$

where it was considered the countable standard model $\mathfrak{M} = (L_{\chi^0}, \in, =)$ of some finite part of the theory $ZF + V = L + \exists k \text{ (} k \text{ is weakly inaccessible cardinal)}$.

In the model $(L_k, \in, =)$ matrix α -functions were considered; such function $\alpha S_f^{<\alpha_1}$ is defined on any nonempty set T^{α_1} , which exist for any sufficiently great cardinal $\alpha_1 < k$, $\alpha_1 \in SIN_n$.

It provides the final contradiction: take any SIN_n -cardinal $\alpha_1 > \alpha\delta^*$ limit for $SIN_n \cap \alpha_1$ of the cofinality $cf(\alpha_1) \geq \chi^{*+}$ providing such nonempty set T^{α_1} with properties (i)–(iii) from theorem 2; such set exist due to lemma 6; then the function $\alpha S_f^{<\alpha_1}$ is nonmonotone on this T^{α_1} by theorem 1 and at the same time is monotone on this set by theorem 2. This contradiction ends the main theorem proof.

References

- [1] Kiselev A. A., *Inconsistency of Inaccessibility*.
arXiv: math.LO http://arxiv.org/PS_cache/arxiv/pdf/1110/1110.3461v1.pdf
- [2] Kiselev A. A., *Inaccessibility and Subinaccessibility*, Second edition, enriched and improved, Part I, Belarusian State University, Minsk, 2008.
arXiv: math.LO http://arxiv.org/PS_cache/arxiv/pdf/1010/1010.1956v4.pdf
- [3] Kiselev A. A., *Inaccessibility and Subinaccessibility*, Second edition, enriched and improved, Part II, Belarussian State University, Minsk, 2010.
arXiv: math.LO http://arxiv.org/PS_cache/arxiv/pdf/1011/1011.1447v2.pdf
- [4] Kiselev A. A., On elimination of subinaccessibility, *Red. zh. Vestsi Akad Navuk BSSR, Ser. Fiz-Mat. Navuk*, 1995, Dep. VINITI N1744-B95 (Russian).
- [5] Kiselev A. A., On generation of subinaccessibility, *Red. zh. Vestsi AN Belarusi, Ser. Fiz-Mat. Navuk*, 1997, Dep. VINITI N3800-B97 (Russian).